

A convergence result of the Lagrangian mean curvature flow

Mu-Tao Wang *

Aug. 15, 2005

email: mtwang@math.columbia.edu

Abstract

We prove the mean curvature flow of the graph of a symplectomorphism between Riemann surfaces converges smoothly as time approaches infinity.

1 Introduction

Let Σ_1 and Σ_2 be two homeomorphic compact Riemann surfaces without boundary. We assume Σ_1 and Σ_2 are both equipped with Riemannian metrics of the same constant curvature c , $c = -1, 0$, or 1 . Let ω_1 and ω_2 denote the volume or symplectic forms of Σ_1 and Σ_2 , respectively. The Riemannian product space $\Sigma_1 \times \Sigma_2$ is denoted by M . We take $\omega' = \omega_1 - \omega_2$ to be the Kähler form of M and M becomes a Kähler-Einstein manifold with the Ricci form $Ric = c\omega'$. Let Σ be the graph of a symplectomorphism $f : \Sigma_1 \rightarrow \Sigma_2$, i.e. $f^*\omega_2 = \omega_1$. Σ can be considered as a Lagrangian submanifold with respect to the symplectic form ω' .

The mean curvature flow deforms the initial surface $\Sigma^0 = \Sigma$ in the direction of its mean curvature vector. Denote by Σ^t the time slice of the flow at t . That Σ^t remains a Lagrangian submanifold follows from a result of Smoczyk [9]. The long-time existence and convergence problems of this flow were studied in [10] and [12].

*The author is partially supported by National Science Foundation Grant DMS0104163 and DMS0306049 and an Alfred P. Sloan Research Fellowship.

In [12], the author proved the long time existence of the flow and showed that Σ^t for $t > 0$ remains the graph of a symplectomorphism f_t . When $c = 1$, the author proved the C^∞ convergence as $t \rightarrow \infty$. However, only C^0 convergence was achieved in the case when $c = -1$ or 0 .

Independently, in [10], Smoczyk studied the case when $c = -1$ or 0 assuming an extra angle condition. He discovered a curvature estimate and showed that the second fundamental form is uniformly bounded under this condition, and thus established the long time existence and C^∞ convergence at infinity.

In view of the above results, it is interesting to see whether the C^∞ convergence of the flow does require the angle condition. In this paper, we show this assumption is unnecessary.

Theorem 1.1 *Let (Σ_1, ω_1) and (Σ_2, ω_2) be two homeomorphic compact Riemann surface of the same constant curvature $c = -1, 0$, or 1 . Suppose Σ is the graph of a symplectomorphism $f : \Sigma_1 \rightarrow \Sigma_2$ as a Lagrangian submanifold of $M = (\Sigma_1 \times \Sigma_2, \omega_1 - \omega_2)$ and Σ^t is the mean curvature flow with initial surface $\Sigma^0 = \Sigma$. Then Σ^t remains the graph of a symplectomorphism f_t along the mean curvature flow. The flow exists smoothly for all time and Σ^t converges smoothly to a minimal Lagrangian submanifold as $t \rightarrow \infty$.*

The long time existence part was already proved in [12]. The smooth convergence was established through a new integral estimate (Lemma 3.1) related to the second variation formula. This estimate is most useful when $c = -1$ or 0 . We remark the existence of such minimal Lagrangian submanifold was proved using variational method by Schoen [7] (see also Lee [5]).

The author would like to thank Tom Ilmanen and Andre Neves for useful discussions.

2 Background material

First we recall some formulas from [12]. The restriction of the Kähler form ω' to Σ^t gives a time-dependent function $\eta = *\omega'$. Since Σ^t is Lagrangian, $*\omega' = 2*\omega_1$. $*\omega_1$ is indeed the Jacobian of the projection π_1 from M to Σ_1 when restricted to Σ and $\eta > 0$ if and only if Σ is locally a graph over Σ_1 . η satisfies the following evolution equation:

$$\frac{d}{dt}\eta = \Delta\eta + \eta[2|A|^2 - |H|^2] + c\eta(1 - \eta^2) \quad (2.1)$$

along the mean curvature flow.

Notice that $0 < \eta \leq 1$. By the equation of η and the comparison theorem for parabolic equations, we get

$$\eta(x, t) \geq \frac{\alpha e^{ct}}{\sqrt{1 + \alpha^2 e^{2ct}}} \quad (2.2)$$

where $\alpha > 0$ is given by $\frac{\alpha}{\sqrt{1+\alpha^2}} = \min_{\Sigma_0} \eta$. Therefore $\eta(x, t)$ converges uniformly to 1 when $c = 1$ and is nondecreasing when $c = 0$. In any case, η has a positive lower bound at any finite time and thus Σ_t remains the graph of a symplectomorphism.

Using the fact that the second fundamental form for Lagrangian submanifold is a fully symmetric three tensor, one derives

$$|H|^2 \leq \frac{4}{3}|A|^2.$$

Plug this into (2.1) and we obtain

$$\frac{d}{dt}\eta \geq \Delta\eta + \frac{2}{3}|A|^2\eta + c\eta(1 - \eta^2) \quad (2.3)$$

In [12], we apply blow-up analysis to this equation to show there exists a weak blow-up limit with vanishing $\int |A|^2$. This together with the lower bound of η shows the limit is a flat space and White's regularity theorem [13] implies the blow-up center is a regular point. This proves the long-time existence of the flow.

3 A monotonicity lemma

In this section, we derive a new monotonicity formula. First $|H|^2$ satisfies the following evolution equation:

$$\left(\frac{d}{dt} - \Delta\right)|H|^2 = -2|\nabla H|^2 + 2 \sum_{ij} \left(\sum_k H_k h_{kij}\right)^2 + c(2 - \eta^2)|H|^2 \quad (3.1)$$

where the symmetric three-tensor h_{ijk} is the second fundamental form and $H_k = h_{iik}$, the trace of the second fundamental form, is the component of the mean curvature vector after identifying the tangent bundle and the normal bundle through J .

We remark that both equations (2.1) and (3.1) are derived in Lemma 5.3 of [10] where p in [10] and η are related by $\eta^2 = \frac{4}{p}$ and $S = 2c$.

We claim the following differential inequality is true:

Lemma 3.1

$$\frac{d}{dt} \int_{\Sigma_t} \frac{|H|^2}{\eta} \leq c \int_{\Sigma_t} \frac{|H|^2}{\eta}$$

Proof.

The proof is a direct computation by combining equations (2.1) and (3.1). We compute

$$\begin{aligned} \frac{d}{dt} \frac{|H|^2}{\eta} &= \frac{\eta \Delta |H|^2 - |H|^2 \Delta \eta}{\eta^2} - 2 \frac{|\nabla H|^2}{\eta} \\ &\quad + \frac{2 \sum_{ij} (\sum_k H_k h_{kij})^2 - 2 |H|^2 |A|^2 + |H|^4}{\eta} + c \frac{|H|^2}{\eta}. \end{aligned}$$

Now

$$\Delta \frac{|H|^2}{\eta} = \frac{\eta \Delta |H|^2 - |H|^2 \Delta \eta}{\eta^2} - \frac{2 \eta \nabla \eta (\eta \nabla |H|^2 - |H|^2 \nabla \eta)}{\eta^4}.$$

We plug this into the previous equation and obtain

$$\begin{aligned} \frac{d}{dt} \frac{|H|^2}{\eta} &= \Delta \frac{|H|^2}{\eta} + \frac{2 \eta \nabla \eta (\eta \nabla |H|^2 - |H|^2 \nabla \eta)}{\eta^4} - 2 \frac{|\nabla H|^2}{\eta} \\ &\quad + \frac{2 \sum_{ij} (\sum_k H_k h_{kij})^2 - 2 |H|^2 |A|^2 + |H|^4}{\eta} + c \frac{|H|^2}{\eta}. \end{aligned}$$

Rearranging terms, we arrive at

$$\begin{aligned} \frac{d}{dt} \frac{|H|^2}{\eta} &= \Delta \frac{|H|^2}{\eta} + \frac{4 \eta |H| |\nabla \eta \cdot \nabla |H| - 2 |\nabla \eta|^2 |H|^2 - 2 \eta^2 |\nabla H|^2}{\eta^3} \\ &\quad + \frac{2 \sum_{ij} (\sum_k H_k h_{kij})^2 - 2 |H|^2 |A|^2 + |H|^4}{\eta} + c \frac{|H|^2}{\eta}. \end{aligned}$$

Integrate this identity and we have

$$\begin{aligned} \frac{d}{dt} \int_{\Sigma_t} \frac{|H|^2}{\eta} &= \int_{\Sigma_t} \frac{4\eta|H|\nabla\eta \cdot \nabla|H| - 2|\nabla\eta|^2|H|^2 - 2\eta^2|\nabla H|^2}{\eta^3} \\ &\quad + \int_{\Sigma_t} \frac{2\sum_{ij}(\sum_k H_k h_{kij})^2 - 2|H|^2|A|^2}{\eta} + c \int_{\Sigma_t} \frac{|H|^2}{\eta}. \end{aligned}$$

We use $|\nabla|H|| \leq |\nabla H|$ in the first summand on the right hand side and complete the square

$$\frac{4\eta|H|\nabla\eta \cdot \nabla|H| - 2|\nabla\eta|^2|H|^2 - 2\eta^2|\nabla|H||^2}{\eta^3} = -2\frac{|\nabla\eta|H| - \eta\nabla|H||^2}{\eta^3}.$$

At last, we apply Cauchy-Schwarz inequality to the second summand and the differential inequality is proved. $\square$

4 Proof of the theorem

The smooth convergence in the case when $c = 1$, i.e. when Σ_1 and Σ_2 are both standard S^2 , was proved in [12].

We prove the C^∞ convergence in the case $c = 0$ and $c = -1$ in the following. By the general convergence theorem of Simon [8], it suffices to show $|A|^2$ is bounded independent of time.

In the case when $c = 0$, by (2.2), η has a positive lower bound. We have

$$\int_{\Sigma_t} |H|^2 \leq \int_{\Sigma_t} \frac{|H|^2}{\eta} \leq K_1 \int_{\Sigma_t} |H|^2 \quad (4.1)$$

for some constant K_1 .

Since $\int_0^\infty \int_{\Sigma_t} |H|^2 < \infty$, there exists a subsequence t_i such that $\int_{\Sigma_{t_i}} |H|^2 \rightarrow 0$ and thus $\int_{\Sigma_{t_i}} \frac{|H|^2}{\eta} \rightarrow 0$ as well. Because $\int_{\Sigma_t} \frac{|H|^2}{\eta}$ is non-increasing, this implies $\int_{\Sigma_t} \frac{|H|^2}{\eta} \rightarrow 0$ for the continuous parameter t as it approaches ∞ . Together with (4.1) this implies $\int_{\Sigma_t} |H|^2 \rightarrow 0$ as $t \rightarrow \infty$. By the Gauss formula, $\int_{\Sigma_t} |A|^2 = \int_{\Sigma_t} |H|^2 \rightarrow 0$. The ϵ regularity theorem in [4] (see also [2]) implies $\sup_{\Sigma_t} |A|^2$ is uniformly bounded.

In the case when $c = -1$, we have

$$\frac{d}{dt} \int_{\Sigma_t} \frac{|H|^2}{\eta} \leq - \int_{\Sigma_t} \frac{|H|^2}{\eta}$$

or

$$\int_{\Sigma_t} \frac{|H|^2}{\eta} \leq K_2 e^{-t}$$

for some constant K_2 ,

Since $\eta \leq 1$, we have

$$\int_{\Sigma_t} |H|^2 \leq K_2 e^{-t}.$$

This implies $\Sigma_t \rightarrow \Sigma_\infty$ in Radon measure. Indeed, for any function ϕ on M with compact support, it is easy to see

$$\frac{d}{dt} \int_{\Sigma_t} \phi = \int_{\Sigma_t} \nabla^M \phi \cdot H + \int_{\Sigma_t} \phi |H|^2,$$

and thus

$$\int_{\Sigma_t} \phi \rightarrow \int_{\Sigma_\infty} \phi.$$

exponentially. Also the limit measure Σ_∞ is unique.

The argument in [12] shows the limit Σ_∞ is smooth. It seems one can adapt the proof of the local regularity theorem of Ecker (Theorem 5.3) [2] or the original local regularity theorem of Brakke [1] to get the uniform bound on second fundamental form. This does require versions of these theorem in a general ambient Riemannian manifold.

We circumvent this step by quoting a theorem in minimal surfaces. Suppose the second fundamental form is unbounded. The blow-up procedure in Proposition 3.1 of [12] produces a limiting flow that exists on $(-\infty, \infty)$. The flow has uniformly bounded second fundamental form $A(x, t)$ and $|A|(0, 0) = 1$. It is not hard to see each slice is the graph of an area-preserving map from $\mathbb{R}^2$ to $\mathbb{R}^2$. Since $\int_{\Sigma_t} |H|^2 \leq K_2 e^{-t}$, the limiting flow will satisfies

$$\int |H|^2 \equiv 0$$

Therefore, we obtain a minimal area-preserving map. A result of Ni [6] generalizing Schoen's theorem [7] shows this is a linear diffeomorphism. This contradicts to the fact that $|A|(0, 0) = 1$.

References

- [1] K. A. Brakke, *The motion of a surface by its mean curvature*. Mathematical Notes, 20. Princeton University Press, Princeton, N.J., 1978.
- [2] K. Ecker, *Regularity theory for mean curvature flow*. Progress in Nonlinear Differential Equations and their Applications, 57. Birkhuser Boston, Inc., Boston, MA, 2004.
- [3] C. J. Earle and J. Eells, *The diffeomorphism group of a compact Riemann surface*. Bull. Amer. Math. Soc. 73 1967 557–559.
- [4] T. Ilmanen, *Singularities of mean curvature flow of surfaces*. preprint, 1997.
- [5] Y.-I. Lee, *Lagrangian minimal surfaces in Kähler-Einstein surfaces of negative scalar curvature*. Comm. Anal. Geom. 2 (1994), no. 4, 579–592.
- [6] L. Ni, *A Bernstein type theorem for minimal volume preserving maps*. Proc. Amer. Math. Soc. 130 (2002), no. 4, 1207–1210.
- [7] R. Schoen, *The role of harmonic mappings in rigidity and deformation problems*. Complex geometry (Osaka, 1990), 179–200, Lecture Notes in Pure and Appl. Math., 143, Dekker, New York, 1993.
- [8] L. Simon, *Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems*. Ann. of Math. (2) 118 (1983), no. 3, 525–571.
- [9] K. Smoczyk, *Der Lagrangesche mittlere Krümmungsfluß (The Lagrangian mean curvature flow)*. Habilitation thesis (English with German preface), University of Leipzig, Germany (1999), 102 pages.
- [10] K. Smoczyk, *Angle theorems for the Lagrangian mean curvature flow*. Math. Z. 240 (2002), no. 4, 849–883.
- [11] M.-T. Wang, *Mean curvature flow of surfaces in Einstein four-Manifolds*. J. Differential Geom. 57 (2001), no. 2, 301–338.
- [12] M.-T. Wang, *Deforming area preserving diffeomorphism of surfaces by mean curvature flow*. Math. Res. Lett. 8 (2001), no.5-6, 651–662.

- [13] B. White, *A local regularity theorem for classical mean curvature flow*. preprint, 1999, revised 2002.